

Bayer Healthcare LLC.

-PE Product Labels –

-continued















RESTRICTED



Drug Facts (continued)

- **Warnings:** Do not use if you are pregnant or breastfeeding. Do not use if you are taking any other drugs that contain aspirin, salicylates, or acetaminophen.
- **Directions:** See the back of the box for directions.
- **Other information:** Do not use if you are allergic to aspirin, salicylates, or acetaminophen.
- **Questions or comments?** See the back of the box for more information.

Alka-Seltzer PLUS Cold

POWERFAST RIZZ NIGHT

SEVERE Cold

LEMON

20 EFFERVESENT TABLETS

Parents: Learn about how medicine works

86925443

1650056752

Drug Facts (continued)

Warnings: Do not use if you are pregnant or breastfeeding. Do not use if you are taking any other drugs that contain aspirin, salicylates, or acetaminophen.

Directions: See the back of the box for directions.

Other information: Do not use if you are allergic to aspirin, salicylates, or acetaminophen.

Questions or comments? See the back of the box for more information.

INSIDE "LIFT HERE" FLAP

Drug Facts (continued)	
<ul style="list-style-type: none"> • you have • asthma • cough with excessive phlegm (mucus) • difficulty urinating due to enlargement of the prostate gland • persistent or chronic cough such as occurs with smoking, asthma, or emphysema • a sodium restricted diet 	<ul style="list-style-type: none"> • thyroid disease • diabetes
Ask a doctor or pharmacist before use if you are	
<ul style="list-style-type: none"> • taking a prescription drug for • gout • diabetes • arthritis 	
When using this product do not exceed recommended dosage	
Stop use and ask a doctor if	
<ul style="list-style-type: none"> • an allergic reaction occurs. Seek medical help right away. • you experience any of the following signs of stomach bleeding: <ul style="list-style-type: none"> • feel faint • vomit blood • have bloody or black stools • have stomach pain that does not get better • pain, cough, or nasal congestion gets worse or lasts more than 7 days • fever gets worse or lasts more than 3 days • redness or swelling is present • new symptoms occur • ringing in the ears or a loss of hearing occurs • cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition. • nervousness, dizziness, or sleeplessness occurs 	
If pregnant or breast-feeding, ask a health professional before use. It is especially important not to use aspirin during the last 3 months of pregnancy unless definitely directed to do so by a doctor because it may cause problems in the unborn child or complications during delivery.	

	
Drug Facts (continued)	
Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away.	
Directions	
<ul style="list-style-type: none"> • adults and children 12 years and over: take 2 tablets fully dissolved in 4 oz of water every 4 hours. Do not exceed 8 tablets in 24 hours or as directed by a doctor. • children under 12 years: do not use. 	
Other information	
<ul style="list-style-type: none"> • each tablet contains: potassium 80 mg, sodium 356 mg • store at room temperature. Avoid excessive heat. 	
Inactive ingredients anhydrous citric acid, calcium silicate, dimethylsiloxane, FD&C red #40, FD&C yellow #6, flavor, mannitol, potassium bicarbonate, povidone, sodium bicarbonate, sucralose.	
Questions or comments? 1-800-966-0369 (Mon-Fri 9AM - 5PM EST)	
	
POWERFAST Fizz™	
NON-DROWSY	
PARENTS:	
Learn about teen medicine abuse	
www.StopMedicineAbuse.org	
19344	
	

Do not use if pouch is torn or open.

Alka-Seltzer PLUS

Cold

CITRUS

POWERFAST Fizz™

NON-DROWSY

Expiry Date: Lot:

D007-14.15

Alka-Seltzer PLUS

SEVERE Cold

CITRUS

POWERFAST Fizz™ **NEW IMPROVED FLAVOR**

NON-DROWSY

ASPIRIN (NSAID) / Pain Reliever-Fever Reducer
Dextromethorphan HBr / Cough Suppressant
Phenylephrine Bitartrate / Nasal Decongestant

- Fever + Body Ache
- Nasal Congestion
- Sinus Pressure
- Sore Throat
- Cough

20 EFFERVESCENT TABLETS

Drug Facts

Active ingredients (in each tablet)

Aspirin 325 mg (NSAID)* Pain reliever/fever reducer
Dextromethorphan hydrobromide 10 mg Cough suppressant
Phenylephrine bitartrate 7.8 mg Nasal decongestant
*nonsteroidal anti-inflammatory drug

Purposes

Uses

- temporarily relieves these symptoms due to a cold with cough:
 - minor aches and pains
 - cough
 - nasal and sinus congestion
 - sore throat
 - temporarily reduces fever
- headache

Warnings

Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product. When using this product, if changes in behavior with nausea and vomiting occur, consult a doctor because these symptoms could be an early sign of Reye's syndrome, a rare but serious illness.

Allergy alert: Aspirin may cause a severe allergic reaction which may include:

- hives
- asthma (wheezing)
- facial swelling
- shock

Drug Facts (continued)

Stomach bleeding warning: This product contains an NSAID, which may cause severe stomach bleeding. The chance is higher if you:

- are age 60 or older
- have had stomach ulcers or bleeding problems
- take a blood thinning (anticoagulant) or steroid drug
- take other drugs containing prescription or nonprescription NSAIDs (aspirin, ibuprofen, naproxen, or others)
- have 3 or more alcoholic drinks every day while using this product
- take more or for a longer time than directed

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- if you are allergic to aspirin or any other pain reliever/fever reducer
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- if you have ever had an allergic reaction to this product or any of its ingredients
- in children under 12 years of age

Ask a doctor before use if

- stomach bleeding warning applies to you
- you have a history of stomach problems, such as heartburn
- you have high blood pressure, heart disease, liver cirrhosis, or kidney disease
- you are taking a diuretic

LIFT HERE

87139344

Dist. by: Bayer HealthCare LLC
Whippany, NJ 07981
Bayer (reg'd), the Bayer Cross (reg'd),
Alka-Seltzer Plus®, and PowerFast Fizz
are trademarks of Bayer.

Bayer

Product of Mexico













Alka-Seltzer Plus® Severe Cough, Mucus & Congestion Day Liquid Gels

Drug Facts

Active ingredients (in each capsule)

Purposes

Acetaminophen 250 mg.....	Pain reliever/fever reducer
Dextromethorphan hydrobromide 10 mg.....	Cough suppressant
Guaifenesin 200 mg.....	Expectorant
Phenylephrine hydrochloride 5 mg.....	Nasal decongestant

Uses

- helps loosen phlegm (mucus) and thin bronchial secretions to rid the bronchial passageways of bothersome mucus and make coughs more productive
- temporarily relieves these symptoms due to a cold or flu:
 - nasal congestion
 - sinus congestion and pressure
 - minor aches and pains
 - headache
 - cough
 - sore throat
- temporarily reduces fever

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take

Alka-Seltzer Plus® Severe Cough, Mucus & Congestion Night Liquid Gels

Drug Facts

Active ingredients (in each capsule)

Purposes

Acetaminophen 325 mg.....	Pain reliever/fever reducer
Dextromethorphan hydrobromide 10 mg.....	Cough suppressant
Doxylamine succinate 6.25 mg.....	Antihistamine
Phenylephrine hydrochloride 5 mg.....	Nasal decongestant

Uses

- temporarily relieves these symptoms due to a cold or flu:
 - minor aches and pains
 - headache
 - nasal and sinus congestion
 - cough
 - sore throat
 - runny nose
 - sneezing
- temporarily reduces fever

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen

- more than 4,000 mg of acetaminophen in 24 hours
- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

LIFT HERE

- 3 or more alcoholic drinks every day while using this product
- Allergy alert:** Acetaminophen may cause severe skin or severe allergic reactions. Symptoms may include:
- skin reddening
 - blisters
 - rash
 - hives
 - facial swelling
 - asthma (wheezing)
 - shock
- If a skin or general allergic reaction occurs, stop use and seek medical help right away.

Cut Edge

8 1/2"

Alka-Seltzer Plus® Severe Cough, Mucus & Congestion Day Liquid Gels

Drug Facts (continued)

Allergy alert: Acetaminophen may cause severe skin or severe allergic reactions. Symptoms may include:

- skin reddening
- blisters
- rash
- hives
- facial swelling
- asthma (wheezing)
- shock

If a skin or general allergic reaction occurs, stop use and seek medical help right away.

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or non-prescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- if you have ever had an allergic reaction to this product or any of its ingredients
- in children under 12 years of age

Ask a doctor before use if you have

- liver disease
- heart disease
- high blood pressure
- thyroid disease
- diabetes
- difficulty in urination due to enlargement of the prostate gland
- persistent or chronic cough such as occurs with smoking, asthma, chronic bronchitis or emphysema
- cough with excessive phlegm (mucus)

Alka-Seltzer Plus® Severe Cough, Mucus & Congestion Night Liquid Gels

Drug Facts (continued)

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use to sedate children.

Do not use

- with any other drug containing acetaminophen (prescription or non-prescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- if you have ever had an allergic reaction to this product or any of its ingredients
- in children under 12 years of age

Ask a doctor before use if you have

- liver disease
- heart disease
- high blood pressure
- thyroid disease
- diabetes
- glaucoma
- cough with excessive phlegm (mucus)
- a breathing problem such as emphysema or chronic bronchitis
- difficulty in urination due to enlargement of the prostate gland
- persistent or chronic cough such as occurs with smoking, asthma, or emphysema


Ask a doctor or pharmacist before use if you are

- taking the blood thinning drug warfarin
- taking sedatives or tranquilizers

PARENTS:

Learn about teen medicine abuse

www.StopMedicineAbuse.org



Alka-Seltzer Plus® Severe Cough, Mucus & Congestion Day Liquid Gels

Drug Facts (continued)

Ask a doctor or pharmacist before use if you are taking the blood thinning drug warfarin

When using this product do not exceed recommended dosage

Stop use and ask a doctor if

- pain, cough, or nasal congestion gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with fever, rash or headache that lasts. These could be signs of a serious condition.
- nervousness, dizziness, or sleeplessness occurs

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- do not take more than the recommended dose
- adults and children 12 years and over: take 2 capsules with water every 4 hours. Do not exceed 10 capsules in 24 hours or as directed by a doctor.
- children under 12 years: do not use

Alka-Seltzer Plus® Severe Cough, Mucus & Congestion Night Liquid Gels

Drug Facts (continued)

When using this product

- do not exceed recommended dosage
- may cause marked drowsiness
- avoid alcoholic drinks
- alcohol, sedatives, and tranquilizers may increase drowsiness
- be careful when driving a motor vehicle or operating machinery
- excitability may occur, especially in children

Stop use and ask a doctor if

- pain, cough, or nasal congestion gets worse or lasts more than 7 days
- fever gets worse or lasts more than 3 days
- redness or swelling is present
- new symptoms occur
- cough comes back or occurs with rash or headache that lasts. These could be signs of a serious condition.
- nervousness, dizziness, or sleeplessness occurs

If pregnant or breast-feeding, ask a health professional before use.

Keep out of reach of children. In case of overdose, get medical help or contact a Poison Control Center right away. Quick medical attention is critical for adults as well as for children even if you do not notice any signs or symptoms.

Directions

- do not take more than the recommended dose
- adults and children 12 years and over: take 2 capsules with water every 4 hours. Do not exceed 10 capsules in 24 hours or as directed by a doctor.
- children under 12 years: do not use

Alka-Seltzer Plus® Severe Cough, Mucus & Congestion Day Liquid Gels

Drug Facts (continued)

Other information

- store at 15-25°C (59-77°F)

Inactive ingredients FD&C red #40, gelatin, glycerin, polyethylene glycol, povidone, propylene glycol, purified water, shellac, sodium hydroxide, sorbitol-sorbitan solution, titanium dioxide

Questions or comments? 1-800-986-0369
(Mon-Fri 9AM - 5PM EST)

Alka-Seltzer Plus® Severe Cough, Mucus & Congestion Night Liquid Gels

Drug Facts (continued)

Other information

- store at 15-25°C (59-77°F)

Inactive ingredients FD&C blue #1, D&C yellow #10, gelatin, glycerin, polyethylene glycol, povidone, propylene glycol, purified water, shellac, sodium hydroxide, sorbitol-sorbitan solution, titanium dioxide

Questions or comments? 1-800-986-0369
(Mon-Fri 9AM - 5PM EST)

Made in Spain

© 2017 Bayer

Dist. by: Bayer HealthCare LLC
Whippany, NJ 07981
Bayer, the Bayer Cross and
Alka-Seltzer Plus are registered
trademarks of Bayer.

85737554

Do not use if blister is cut or broken.

DOES NOT CONTAIN ASPIRIN

Bayer

LOT/EXP



**LOT & EXP
CODE AREA**



3/4

1/2





Lot no:
Exp:



LIFT HERE

Drug Facts (continued)

Sore throat warning: If sore throat is severe, persists for more than 2 days, is accompanied or followed by fever, headache, rash, nausea, or vomiting, consult a doctor promptly.

Do not use

- with any other drug containing acetaminophen (prescription or nonprescription). If you are not sure whether a drug contains acetaminophen, ask a doctor or pharmacist.
- if you are now taking a prescription monoamine oxidase inhibitor (MAOI) (certain drugs for depression, psychiatric, or emotional conditions, or Parkinson's disease), or for 2 weeks after stopping the MAOI drug. If you do not know if your prescription drug contains an MAOI, ask a doctor or pharmacist before taking this product.
- if you have ever had an allergic reaction to this product or any of its ingredients
- in children under 12 years of age

Ask a doctor before use if you have

- liver disease
- heart disease
- high blood pressure
- thyroid disease
- diabetes
- cough with excessive phlegm (mucus)
- difficulty in urination due to enlargement of the prostate gland
- persistent or chronic cough such as occurs with smoking, asthma, or emphysema

Ask a doctor or pharmacist before use if you are taking the

Blood thinning drug warfarin

When using this product do not exceed recommended dosage

Drug Facts

Active ingredients (in each capsule)

Acetaminophen 325 mg Pain reliever/fever reducer

Dextromethorphan hydrobromide 10 mg Cough suppressant

Phenylephrine hydrochloride 5 mg Nasal decongestant

Uses

- temporarily relieves these symptoms due to a cold or flu:
- minor aches and pains
- headache
- cough
- nasal and sinus congestion
- temporarily reduces fever

Warnings

Liver warning: This product contains acetaminophen. Severe liver damage may occur if you take more than 4,000 mg of acetaminophen in 24 hours

- with other drugs containing acetaminophen
- 3 or more alcoholic drinks every day while using this product

Allergy alert: Acetaminophen may cause severe skin or severe allergic reactions. Symptoms may include:

- skin reddening
- skin swelling
- hives
- rash
- blisters
- asthma (wheezing)
- shock

If a skin or general allergic reaction occurs, stop use and seek medical help right away.

Alka-Seltzer PLUS

MAXIMUM STRENGTH

Cold & Flu

DAY NON-DROWSY

Alka-Seltzer PLUS

MAXIMUM STRENGTH

Cold & Flu

DAY NON-DROWSY

ACETAMINOPHEN / Pain Reliever-Fever Reducer
Dextromethorphan Hydrobromide / Cough Suppressant
Phenylephrine HCl / Nasal Decongestant

- ▶ Cough
- ▶ Nasal Congestion
- ▶ Headache & Body Ache
- ▶ Sore Throat
- ▶ Sinus Pressure




20 LIQUID GELS (Liquid Filled Capsules)



6256

6256



<p>Drug Facts (continued)</p> <ul style="list-style-type: none"> ● redness or swelling is present ● new symptoms occur ● ringing in the ears or loss of hearing occurs ● cough comes back or occurs with rash or headache that lasts. There could be signs of a serious condition. ● nervousness, dizziness, or sleeplessness occurs 	<p>Drug Facts (continued)</p> <ul style="list-style-type: none"> ● operating machinery ● excitability may occur, especially in children Stop use and ask a doctor if <ul style="list-style-type: none"> ● an allergic reaction occurs. Seek medical help right away. ● you experience any of the following signs of stomach bleeding: <ul style="list-style-type: none"> ● feel faint ● vomit blood ● have black or black stools 	<p>Drug Facts (continued)</p> <ul style="list-style-type: none"> ● minor aches and pains ● headache ● nasal and sinus congestion ● temporarily reduces fever <p>Warnings</p> <p>Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product.</p>	<p>Drug Facts (continued)</p> <ul style="list-style-type: none"> ● minor aches and pains ● headache ● sinus congestion and pressure ● cough ● sore throat ● temporarily reduces fever <p>Warnings</p> <p>Reye's syndrome: Children and teenagers who have or are recovering from chicken pox or flu-like symptoms should not use this product.</p>	<p>UPPER PANEL</p>
--	---	--	---	---------------------------